

Electric Characteristics of Rotational States positive parity in isotopes $^{170,172,174}\text{Yb}$

P.N.Usmanov*

Institute of Nuclear Physics, Academy of Science of Uzbekistan, 100214, Ulugbek, Tashkent, Uzbekistan

A.A.Okhunov†

*Department of Science in Engineering, KOE, International Islamic University Malaya,
P.O Box 10, 50728 Kuala Lumpur, Malaysia and*

Institute of Nuclear Physics, Academy of Science of Uzbekistan, 100214, Ulugbek, Tashkent, Uzbekistan

H. Abu Kassim‡

Department of Physics, University of Malaya, 50603 Kuala Lumpur, Malaysia

U.S. Salikhbaev§

*Institute of Nuclear Physics, Academy of Science of Uzbekistan, 100214, Ulugbek, Tashkent, Uzbekistan
(Dated: May 29, 2015)*

Accounting for Coriolis mixing of experimentally known rotational bands with $K^\pi < 3^+$, non-adiabatic effects in energy and electric characteristics of excited states are investigated, within phenomenological model.

The energy and wave function structure of excited states are calculated. The finding reveals that the bands mixing has been found to have considerable impact on the wave function of low-lying states 0^+ and 2^+ bands.

In addition, the probabilities of $E2$ - transitions have been calculated. The values from calculations of $B(E2)$ - transitions from 0_2^+ , 0_3^+ , 2_1^+ , and 2_2^+ bands are compared with the experimental data.

PACS numbers: 21.10.-k, 21.10.Re, 21.10.Ky, 21.10.Hw

I. INTRODUCTION

Despite the fact that the structure of deformed nuclei and nature of low excited levels have been substantially studies over more than four decades, this still occupies a central part of today's research [1]-[3].

An extensive research interest in the properties of deformed nuclei has risen in recent years with the exploration of a new collective isovector magnetic dipole mode [4, 5]. The measured values of excited energy of magnetic mode are found to be not so high in an excited spectrum, and consideration of mixing with low-lying exciting states appear to lead to an interesting physical phenomena [6, 7].

The nuclei $^{170,172,174}\text{Yb}$ have been well studied. It is important to note that these are investigated in a number of ways such as radioactive decay of $^{170,172,174}\text{Lu}$, and different nuclear reactions. In these isotopes, many 1^+ states and $K^\pi = 0^+, 2^+$ bands have been observed. For instance, the excited energy $K^\pi = 0^+$ and 2^+ it rises with the increase in number of neutrons (see Figure 1).

The values of probability of $B(E2)$ with low-lying levels of $K^\pi = 0^+, 2^+$ bands, and also Rasmusson's parameter value $X_I(E0/E2)$, dimensionless units matrix




FIG. 1: (Color online) The energy of $I = 2$ states of $K^\pi = 0_2^+, 0_3^+, 0_4^+, 0_5^+, 2_1^+$ and 2_2^+ bands in isotopes $^{170,172,174}\text{Yb}$.

element $\rho(E0)$ for the $E0$ - transition and a multipole mixture coefficients $\delta(E2/M1)$ are defined experimentally [8]-[10].

Numerous conducted experiments on defining spectroscopic characteristics of low-lying exciting states, particular $K^\pi = 1^+$ in deformed nuclei [5], have motivated the further theoretical investigations. In this case, investigations influence of $K^\pi = 1^+$ states to the properties of low-lying levels is actual.

*Electronic address: usmanov1956.56@mail.ru

†Electronic address: Corresponding author: aaokhunov@gmail.com;
URL: <http://www.iium.edu.my>

‡Electronic address: hasank@um.edu.my

§Electronic address: salikhbaev@inp.uz

Present paper focuses on low-lying states of positive parity of isotopes $^{170,172,174}\text{Yb}$. The calculation are conducted by utilizing a phenomenological model [6] which accounts Coriolis mixture all of the experimentally known low-lying rotational bands states with $K^\pi < 3^+$.

Experimentally observed K^- forbidden transitions as well as nonadiabaticities of energy and in ratios of $E2^-$ transitions can be explained by Coriolis mixture states.

II. THE MODEL

To analyze the properties of low-lying positive parity states in Yb isotopes, the phenomenological model of [6] is exploited. This model takes into account the mixing of states of the $K^\pi = 0^+, 2^+$ and 1^+ bands. The Hamiltonian model is

$$H = H_{rot}(I^2) + H_{K,K'} \quad (1)$$

$$H_{K',K}^\sigma(I) = \omega_K \delta_{K,K'} - \omega_{rot}(I)(j_x)_{K,K'} \zeta(I, K) \delta_{K,K' \pm 1} \quad (2)$$

where ω_K —bandhead energy of rotational band, $\omega_{rot}(I)$ —an angular frequency of rotational nucleus, $(j_x)_{K,K'}$ —matrix elements which describe Coriolis mixture between rotational bands and

$$\zeta(I, 0) = 1 \quad \zeta(I, 2) = \left[1 - \frac{2}{I(I+1)} \right]^{\frac{1}{2}}$$

The eigenfunction of Hamiltonian model (1) is

$$\begin{aligned} |IMK\rangle &= \sqrt{\frac{2I+1}{16\pi^2}} \left\{ \sqrt{2} \Psi_{gr,K}^I D_{MK}^I(\theta) \right. \\ &+ \sum_{K'} \frac{\Psi_{K',K}^I}{\sqrt{1+\delta_{K',0}}} [D_{M,K'}^I(\theta) b_{K'}^+ \\ &+ (-1)^{I+K'} D_{M,-K'}^I(\theta) b_{-K'}^+] \left. \right\} |0\rangle \end{aligned} \quad (3)$$

here $\Psi_{K',K}^I$ is the amplitude of mixture of basis states.

The rotational part of Hamiltonian (1) $H_{rot}(I)$ is diagonal by wave functions (3). Note that $H_{rot}(I)$ is determined by exploiting Harris parameterization for energy and angular momentum [11]

$$E_{rot}(I) = \frac{1}{2} \mathfrak{S}_0 \omega_{rot}^2(I) + \frac{3}{4} \mathfrak{S}_1 \omega_{rot}^4(I) \quad (4)$$

$$[I(I+1)]^{1/2} = \mathfrak{S}_0 \omega_{rot}(I) + \mathfrak{S}_1 \omega_{rot}^3(I) \quad (5)$$

where $\mathfrak{S}_0$ and $\mathfrak{S}_1$ —are the inertia parameters of the rotational core.

The rotational frequency of the core $\omega_{rot}(I)$ is found by solving cubic equation (5). This equation has two

imaginary roots and one real root. The real root is as follows

$$\begin{aligned} \omega_{rot}(I) &= \left\{ \frac{\tilde{I}}{2\mathfrak{S}_1} + \left(\left(\frac{\tilde{I}}{2\mathfrak{S}_1} \right)^2 + \left(\frac{\mathfrak{S}_0}{3\mathfrak{S}_1} \right)^3 \right)^{\frac{1}{2}} \right\}^{\frac{1}{3}} \\ &+ \left\{ \frac{\tilde{I}}{2\mathfrak{S}_1} - \left(\left(\frac{\tilde{I}}{2\mathfrak{S}_1} \right)^2 + \left(\frac{\mathfrak{S}_0}{3\mathfrak{S}_1} \right)^3 \right)^{\frac{1}{2}} \right\}^{\frac{1}{3}} \end{aligned} \quad (6)$$

where $\tilde{I} = \sqrt{I(I+1)}$. Equation (6) gives $\omega_{rot}(I)$ at the given spin I of the core.

Solving the Shrödinger equation

$$H_{K,K'}^\sigma \Psi_{K,K'}^I = \mathcal{E}_K(I) \Psi_{K,K'}^I. \quad (7)$$

we define eigne function and energy of a Hamiltonian. The total energy of state is defined by

$$E_K(I) = E_{rot}(I) + \mathcal{E}_K(I) \quad (8)$$

A. Energy spectra and structures of the states

The calculations have been carried out for the isotopes $^{170,172,174}\text{Yb}$. All experimentally known rotational bands of positive parity with $K^\pi < 3^+$ have been included in basis Hamiltonian states.

The experiment suggests that $m = 5$ band with $K^\pi = 0_m^+$, one band $\ell = 1$ with $K^\pi = 2_\ell^+$, and $\nu = 19$ with $K^\pi = 1_\nu^+$ states in ^{170}Yb [8]. These all $n = m + \ell + \nu = 25$ rotational bands have been included in the basis states of Hamiltonian (1). For the isotopes $^{172,174}\text{Yb}$, basis states of Hamiltonian include $n = 15$ ($m = 5$, $\ell = 2$ and $\nu = 8$) and $n = 22$ ($m = 5$, $\ell = 2$ and $\nu = 15$), correspondingly [5, 9, 10].

The parameters of inertia $\mathfrak{S}_0$ and $\mathfrak{S}_1$ are estimated by exploiting Harris parameterization (4), and using the experimental data for energy up to spin $I \leq 8\hbar$ for ground band [12].

The Hamiltonian (2) has transformational properties, that the state (3) can be classified as quantum number — $\sigma = \pm 1$ signature, which imposes restrictions on angular momentum values.

$$(-1)^I \sigma = 1.$$

For the states with negative signature $\sigma = -1$, Hamiltonian (2) has dimension $n = \ell + \nu$, as in bands with $K^\pi = 0_m^+$ the are no condition states with odd spins I . For the states with positive signature $\sigma = +1$, Hamiltonian (2) has dimension $n = m + \ell + \nu$.

The model parameters are described as follows:

a) the bandhead energy ground ω_{0_1} and $K^\pi = 0_m^+$ bands has taken from experiment, as they are not revolted by Coriolis force. Bandhead energy of 1_ν^+ bands

are also defined from an experiment [5, 8]

$$\omega_{1\nu} = E_{1\nu}^{exp.}(I=1) - E_{rot}(I=1);$$

b) matrix elements $(j_x)_{2\ell 1} = (j_x)_{2\ell 1\nu}$ and bandhead energy of 2_ℓ^+ bands $\omega_{2\ell}$ are determined from the most favored experimental and theoretical spectrum of energy states with a negative signature $\sigma = -1$, e.a. for energy

state for even spins I ;

c) the matrix elements $(j_x)_{0_m 1} = (j_x)_{0_m 1\nu}$ defined by the least square method from the best fitted of theoretical energy spectra state with positive signature $\sigma = +1$ with experimental data.

The obtained values of model parameters are presented in Table 1.

TABLE I: Parameters used in the calculations for the isotopes $^{170,172,174}Yb$

A	$(j_x)_{01,1\nu}$	$(j_x)_{02,1}$	$(j_x)_{03,1}$	$(j_x)_{04,1}$	$(j_x)_{05,1}$	$(j_x)_{21,1}$	$(j_x)_{22,1}$
170	0.1864	0.3936	0.6586	0.9081	0.0009	0.7278	—
172	0.2754	0.9777	0.7176	0.11	0.30	0.325	0.21
174	0.185	0.4	0.25	0.15	0.20	0.085	0.1

Note: $(j_x)_{K',K}$ are matrix elements of the Coriolis interactions.

Calculation comparison of energy with experimental values for $^{170,172,174}Yb$ is illustrated in Figures 2,3 and 4, correspondingly.

Apparently, one may see from the Figures that the model qualitatively reproduces experimental energy of rotational states up to energy $3MeV$. However, in high spin values $I \geq 12\hbar$ noticeable deviation has been observed in calculated values of energy and that obtained from experiment. Note that this deviation increases with the growth of angular momentum I . This is probably due to the fact that the influence of rotation on internal nuclei structure has not been considered in this model. In future, we will study electromagnetic properties of low-lying states $I < 10\hbar$.

Amplitude of the states $\Psi_{K',K}$ for $K^\pi = 2_1^+$ and 0_2^+ bands for $^{170,172,174}Yb$, are provided in Figures 5, 6, 7, 8, 9 and 10, respectively. The components which have small values are not illustrated in Figure. Also the components $\Psi_{1\nu,K}$ band are not given except for the first 1_1^+ . The values for others $\Psi_{1\nu,K}$ states are define as follows

$$\Psi_{1\nu,K} = \Psi_{11,K} \frac{\omega_{11} - \omega_K}{\omega_{1\nu} - \omega_K}. \quad (9)$$

From Figures 5 and 7, we can see that $K^\pi = 0_2^+$ and

$K^\pi = 2_1^+$ bands states in ^{170}Yb , $K^\pi = 0_3^+$ and $K^\pi = 2_1^+$ bands in ^{172}Yb are mixed strongly even in low spin values I . It is associated with the close location to each other (see Figure 1). In isotopes $^{170,172}Yb$, considerable deviation in signature of the states $K^\pi = 2_1^+$ band can be observed. This reflects in the values of probability electromagnetic transitions. In this case, description of quantum number K is difficult for these states. Thus, in ^{170}Yb , a number of research works [13] in this context note that the states with $I = 2^+$ (1.1386 MeV) and $I = 2^+$ (1.1454 MeV) $K = 2$ and $K = 0$, respectively. On the other hand, some works [8] document that $K = 0$ and $K = 2$, correspondingly. In case ^{174}Yb , the mixture effect is not so strong.

III. ELECTRIC QUADRUPOLE TRANSITIONS

With the wave functions calculated by solving the Schrödinger equation (7), reduced probabilities of $E2$ -transitions between states $I_i K_i$ and states of ground band $I_f 0_1$ are calculated [6].

$$B(E2; I_i K_i \rightarrow I_f 0_1) = \left\{ \sqrt{\frac{5}{16\pi}} e Q_0 \left[\Psi_{01,01}^{I_f} \Psi_{01,K_i}^{I_i} C_{I_i 0;20}^{I_f 0} + \sum_n \Psi_{K_n,01}^{I_f} \Psi_{K_n,K_i}^{I_i} C_{I_i K_n;20}^{I_f K_n} \right] + \sqrt{2} \left[\Psi_{01,01}^{I_f} \sum_n \frac{(-1)^{K_n} m_{K_n} \Psi_{K_n,K_i}^{I_i}}{\sqrt{1 + \delta_{K_n,0}}} C_{I_i K_n;2-K_n}^{I_f 0} + \Psi_{01,K_i}^{I_i} \sum_n \frac{m_{K_n} \Psi_{K_n,01}^{I_f}}{\sqrt{1 + \delta_{K_n,0}}} C_{I_i 0;2K_n}^{I_f K_n} \right] \right\}^2 \quad (10)$$

here $m_{K_n} = \langle 0_1^+ | \hat{m}(E2) | K_n^+ \rangle$ is matrix elements between intrinsic wave functions of ground (0_1^+) and $K_n^+ = 0_m^+, 2_\ell^+, 1_\nu^+$ bands which has a value obtained from experimental data, Q_0 is nuclear intrinsic quadrupole moment; and $C_{I_i K_i; 2(K_i + K_1)}^{I_f K_f}$ - Clebsch-Gordan coefficient.

For the reduced probabilities of $E2$ - transitions from the $I = 2$ state we have following equation in adiabatic approximation:

$$B^{adia}(E2; 2K_n \rightarrow 00_1) = (2 - \delta_{K_n, 0}) |m_{K_n} C_{2K_n; 2-K_n}^{00}|^2 \quad (11)$$

which allows us to identify m_{K_n} parameter from the experimental data.

The $K^\pi = 0_2^+, 0_3^+$ and 2_1^+ bands are very close to each other in isotopes $^{170,172}\text{Yb}$, which leads to a strong mixing of states even $I = 2$. In this case, the adiabatic approximation (11) becomes inapplicable to determine m_{K_i} .

The magnitude and sign parameters $m_{1_1} = m_{1_\nu}$ and $m_{2_\ell^+}$ are obtained from the best fitted ratio probabilities $R_{IK} = B(E2; IK \rightarrow I + 10_1) / B(E2; IK \rightarrow I - 10_1)$ and odd states of $K^\pi = 2_\ell^+$ and 1_ν^+ bands. In addition, the most favored ratio R_{IK} and even states (for the positive signature $\sigma = +1$) help to identify parameters m_{0_m} .

Table 2 reports m_K parameters which have been used in calculation $E2$ - transitions.

TABLE II: The values of the parameters m_K and the intrinsic quadrupole moment Q_0 , which are used in calculations (in efm^2)

A	m_{0_2}	m_{0_3}	m_{0_4}	m_{0_5}	m_{1_ν}	m_{2_1}	m_{2_2}	$Q_0[13]$
170	2	24	3	8	-5	19	8	780(4)
172	10	1	-6.9	-8	-5	15	-8	791(4)
174	8	1	-6.9	8	-10 ($m_{1_1} = -1.7$)	15	8	782(4)

Table 3 compares reduced probability $E2$ - transitions with existing experimental data [8]-[10], [14]. Moreover, reduced matrix elements of $E2$ - transitions for ^{172}Yb are

provided in Table 4. In a similar vein, these values are also compared with experimental values as well as values found by using other models [15]-[17].

TABLE III: Reduced probability of $E2$ - transitions in the isotopes Yb (e^2fm^4)

A	$I_i K_i \rightarrow I_f K_f$	Exp.	Theory	$I_i K_i \rightarrow I_f K_f$	Exp.	Theory
^{170}Yb	$22_1 \rightarrow 00_1$	151(35)[8]	90	$00_3 \rightarrow 20_1$	60(15)[8]	43
	$\rightarrow 20_1$	269(60)[8]	60	$00_4 \rightarrow 20_1$	567(118)[8]	567
	$\rightarrow 40_1$	27(6)[8]	10			
^{172}Yb	$22_1 \rightarrow 00_1$	75.6(63)[9]; 74.6(57)[14]	82	$00_2 \rightarrow 20_1$	205(60)[9]	100
	$\rightarrow 20_1$	121(12)[14]	130	$20_2 \rightarrow 00_1$	14(1)[9]; 14(1)[14]	13
	$\rightarrow 40_1$	7.3(6)[9]; 6.8(7)[14]	8.6	$\rightarrow 20_1$	45(7)[9]; 52(8)[14]	23
	$42_1 \rightarrow 20_1$	398(284)[9]	15	$\rightarrow 40_1$	142(20)[9]; 140(20)[14]	74
	$\rightarrow 40_1$	739(512)[9]	81	$00_3 \rightarrow 20_1$	0.14(3)[9]	1
	$32_1 \rightarrow 20_1$	152(11)[14]	154	$20_3 \rightarrow 00_1$	0.4(1)[9]; 3.4(2)[14]	3.6
	$\rightarrow 40_1$	79(6)[14]	73	$\rightarrow 20_1$	0.6(4)[9]; 11.9(8)[14]	3.0
	$22_2 \rightarrow 00_1$	20(4)[9]; 32(4)[14]	23	$\rightarrow 40_1$	1.0(1)[14]	1.2
	$\rightarrow 20_1$	31(2)[9]; 51(7)[14]	38	$00_4 \rightarrow 20_1$	>0.25[9]	48
	$\rightarrow 40_1$	3.3(4)[14]	2.2	$20_4 \rightarrow 00_1$	10(6)[9]	12
	$32_2 \rightarrow 20_1$	54(7)[14]	42	$00_5 \rightarrow 20_1$	> 0.27[9]	64
	$\rightarrow 40_1$	22(3)[14]	21	$20_5 \rightarrow 00_1$	19(9)[9]	21
				$\rightarrow 20_1$	> 0.18[9]	23
^{174}Yb	$00_2 \rightarrow 20_1$	81 $^{+64}_{-29}$ [10]	64	$22_1 \rightarrow 20_1$	144(30)[10]	133

It is important to note that our results are obtained consecutively. In the initial step, energy and wave func-

tion the states are computed. Further, by utilizing these

wave functions, reduced probability of $E2^-$ transitions are calculated. From the Table 4, one may gather that

performed calculations within our model provide a better correspondence with experiment data.

TABLE IV: Reduced Matrix Elements of $E2^-$ transitions in ^{172}Yb , calculated within our model which comparison with experimental data [15] and are calculated using the rotational-vibrational model (RVM2) [16] and the IBA-1 model [17] (eb)

$I_i K_i \rightarrow I_f K_f$	Exp.	RVM2	IBA-1	Theory	$I_i K_i \rightarrow I_f K_f$	Exp.	RVM2	IBA-1	Theory
$20_1 \rightarrow 20_1$	$-2.63^{+0.28}_{-0.27}$	-2.92	-2.92	-2.93	$20_1 \rightarrow 00_1$	$2.45^{+0.12}_{-0.12}$	$2.45^a)$	$2.45^a)$	$2.45^a)$
$40_1 \rightarrow 40_1$	$-3.54^{+0.84}_{-0.18}$	-3.73	-3.69	-3.74	$40_1 \rightarrow 20_1$	$3.76^{+0.19}_{-0.19}$	3.93	3.91	3.93
$60_1 \rightarrow 60_1$	$-4.31^{+0.22}_{-0.62}$	-4.43	-4.33	-4.46	$60_1 \rightarrow 40_1$	$5.34^{+0.27}_{-0.27}$	4.97	4.90	4.96
$80_1 \rightarrow 80_1$	$-4.49^{+0.23}_{-0.77}$	-5.05	-4.83	-5.08	$80_1 \rightarrow 60_1$	$5.90^{+0.30}_{-0.30}$	5.80	5.60	5.80
$100_1 \rightarrow 100_1$	$-6.32^{+0.74}_{-0.32}$	-5.60	-5.22	-5.63	$100_1 \rightarrow 80_1$	$6.71^{+0.34}_{-0.34}$	6.54	6.29	6.54
$120_1 \rightarrow 120_1$	$-6.15^{+0.64}_{-0.74}$	-6.08	-5.53	-6.15	$120_1 \rightarrow 100_1$	$7.01^{+0.35}_{-0.35}$	7.19	6.79	7.20
$140_1 \rightarrow 140_1$	—	—	—	6.62	$140_1 \rightarrow 120_1$	$8.12^{+0.63}_{-0.43}$	7.80	7.18	7.81
$22_1 \rightarrow 00_1$	$0.208^{+0.010}_{-0.040}$	0.21	$0.20^a)$	0.203	$00_2 \rightarrow 20_1$	$0.166^{+0.018}_{-0.018}$	0.16	0.27	0.01
$\rightarrow 20_1$	$0.250^{+0.016}_{-0.018}$	0.25	0.31	0.255	$20_2 \rightarrow 00_1$	$0.090^{+0.010}_{-0.040}$	0.16	0.26	0.082
$\rightarrow 40_1$	$0.063^{+0.009}_{-0.004}$	0.062	0.10	0.066	$\rightarrow 20_1$	$-0.162^{+0.071}_{-0.008}$	0.19	-0.31	0.108
$42_1 \rightarrow 20_1$	$0.22^{+0.07}_{-0.05}$	0.20	0.13	0.11	$\rightarrow 40_1$	$0.27^{+0.02}_{-0.08}$	0.26	0.45	0.19
$\rightarrow 40_1$	$0.46^{+0.08}_{-0.13}$	0.38	0.45	0.27	$40_2 \rightarrow 40_1$	—	—	—	0.13
$32_1 \rightarrow 20_1$	0.32(11)	—	—	0.328					
$\rightarrow 40_1$	0.235(6)	—	—	0.226					

^{a)} This matrix element was used to normalize the results of the model calculations.

To evaluate the degree of nonadiabaticity, manifested in the reduced probabilities of $E2^-$ transitions, in Table 5 theoretical ratios R_{IK} has been compared with their adiabatical values $R_{IK}^{adiab.}$ as well as experimental data [8]-[10], [13, 14, 18] which is determined as follows

$$R_{IK} = \frac{I_\gamma(IK \rightarrow I_1 0_1)}{I_\gamma(IK \rightarrow I_2 0_1)} \left(\frac{E_\gamma(IK \rightarrow I_2 0_1)}{E_\gamma(IK \rightarrow I_1 0_1)} \right)^5 \quad (12)$$

In ^{170}Yb , experimental ratio R_{I0_2} for $E2^-$ transitions from states $K^\pi = 0_2^+$ band differ from adiabatic theory considerably (10-40 times). This is associated with mixing 0_2^+ and 2_1^+ bands. An important question can raised in this regard. Why do the ratios R_{I2_1} for transitions from 2_1^+ bands differ not so strongly from adiabatic theory with respect to R_{I0_2} ?

This results can be explained by the fact that the matrix element m_{2_1} is greater about 10 times than that of m_{0_2} (see Table 2). One may see from this comparison that the mixing effect of states low-lying bands plays a crucial role which considerably demonstrates that $E2^-$ transitions even in low values of angular momentum I .

where $I_\gamma(IK \rightarrow I_1 0_1)$ — is intensity and $E_\gamma(IK \rightarrow I_1 0_1)$ — is energy of γ^- transition.

IV. CONCLUSION

In the present work, non-adiabatic effects in energies and electric characteristics of excited states are studied within the phenomenological model which taking into account Coriolis mixing of all experimentally known rotational bands with $K^\pi < 3^+$.

The energy and structure of wave functions of excited states are calculated. And also the reduced probabilities of $E2^-$ transitions is calculated. The ratio of $E2^-$ transitions probability from $K^\pi = 0_m^+$ and 2_ℓ^+ bands are calculated and compared with experimental data which gives the satisfactory result.

If matrix elements of $E2^-$ transitions m_K one of two strongly mixing bands K is less than matrix element of

TABLE V: The ratios of reduced probabilities of $E2$ - transitions $R_{IK} = B(E2; I_i K_i \rightarrow I_f 0_1) / B(E2; I_i K_i \rightarrow I'_f 0_1)$ in isotopes Yb

A	$I_i K_i$	$I_f K_f$	$I'_f K'_f$	Experiments		Theory	Alaga
^{170}Yb	22_1^+	20_1^+	00_1^+	1.77(8)[8]		1.86	1.43
		40_1^+	20_1^+	0.098(11)[8]		0.043	0.050
		40_1^+	20_1^+	0.78(4)[8]		0.75	0.40
		60_1^+	40_1^+	1.39(46)[8]		1.50	0.57
	72_1^+	80_1^+	60_1^+	1.27(24)[8]		2.42	0.67
	20_2^+	20_1^+	00_1^+	1.94(52)[8]		1.1	1.43
		40_1^+	20_1^+	—		1.82	0.91
		60_1^+	40_1^+	3.73(90)[8]		1.67	0.81
		80_1^+	60_1^+	10.7(19)[8]		2.45	0.77
		100_1^+	80_1^+	25.3(85)[8]		29.2	0.74
		120_1^+	100_1^+	29.9(71)[8]		2.0	0.73
	20_3^+	20_1^+	00_1^+	1.81(11)[8]		2.5	1.43
		40_1^+	20_1^+	3.00(15)[8]		2.3	1.80
		40_1^+	00_1^+	5.43(18)[8]		5.75	2.57
		40_1^+	00_1^+	4.0(2)[8]		3.9	2.57
^{172}Yb	22_1^+	20_1^+	00_1^+	1.62(12) [14]	1.71(84) [9]	1.59	1.43
		40_1^+	20_1^+	0.056(5) [14]	0.056(20) [9]	0.066	0.072
		40_1^+	20_1^+	0.52(4) [14]	0.56(3) [9]	0.48	0.40
		40_1^+	20_1^+		3.35(69) [9]	6.0	2.94
	22_2^+	20_1^+	00_1^+	1.69(21) [14]	1.55(30) [9]	1.61	1.43
		40_1^+	20_1^+	0.015(8) [14]	0.064(38) [9]	0.058	0.072
		40_1^+	20_1^+	0.163(56) [14]	0.409(77) [9]	0.504	0.40
		40_1^+	20_1^+		4.11(17) [9]	3.89	2.94
	52_2^+	60_1^+	40_1^+		< 1.10 [9]	0.82	0.57
		20_1^+	00_1^+	3.71(24) [14]	2.88(36) [9]	1.80	1.43
	40_2^+	40_1^+	20_1^+	2.70(38) [14]	2.61(11) [9]	3.20	1.80
		40_1^+	20_1^+		6.78(1.36) [9]	1.56	0.91
^{174}Yb	22_1^+	20_1^+	00_1^+	> 0.49[13, 18]	2.4(5)[10]	1.59	1.43
		40_1^+	20_1^+	0.167(75)[13, 18]	0.256(92)[10]	0.055	0.072
		40_1^+	20_1^+	> 0.325[13, 18]	0.67(13)[10]	0.49	0.40
		40_1^+	20_1^+	4.83(5)[13, 18]	4.77(63)[10]	3.75	2.94
	20_2^+	60_1^+	40_1^+	≤ 0.14 [13, 18]		0.09	0.086
		20_1^+	00_1^+	> 8.8[13, 18]	> 9.7[10]	2.63	1.43
	40_2^+	40_1^+	20_1^+	3.45(5)[13, 18]	2.9(2)[10]	6.1	1.80
		40_1^+	20_1^+	7.82(26)[13, 18]	11.8(2.5)[10]	6.21	0.91
		60_1^+	40_1^+	1.50(22)[13, 18]	0.59(7)[10]	11.3	1.75
	20_4^+	20_1^+	00_1^+	> 1.7[13, 18]	1.99(27)[10]	1.54	1.43
		40_1^+	20_1^+	1.24(16)[13, 18]	1.12(12)[10]	2.11	1.80

$m_{K'}$ of K' ($m_K < m_{K'}$), then the difference in the ratio R_{IK} for the first band K from Alaga rule is bigger than the difference in $R_{IK'}$ from Alaga rule. In other words, if $m_K < m_{K'}$, nonadiabaticity in the ratio R_{IK} is stronger than that of $R_{IK'}$.

Acknowledgments

This work has been financial supported by the MOHE, Fundamental Research Grant Scheme FRGS13-074-0315.

We thank the Islamic Development Bank (IDB) for supporting this work under scholarships 37/IRQ/P30.

[1] A.Bohr and B.R.Mottelson, "Nuclear Structure Vol. **1.2**" Benjamin, New York, (1997).

[2] V.G.Soloviev "Theory of difficult atomic nuclei" // M.

- Nauk, (1974).
- [3] D.G.Burke, V.G.Soloviev, A.V.Sushkov and N.Yu.Shirikova // Nucl. Phys. **A656**, 287 (1999).
 - [4] D.Bohle, G.Kuchler, A.Richter and W.Steffen // Phys.Lett. **B4,5**(148), 411 (1984).
 - [5] A.Zilges, P. von Brentano, C.Wesselborg and et al. // Nuc.Phys. **A507**, p.399 (1990).
 - [6] Ph.N.Usmanov, I.N.Mikhailov.// Phys. Part. Nucl. Lett. **28**, 348 (1997) [Fiz.Elem.Chastits At. Yadra **28**, p.887 (1997)].
 - [7] Ph.N.Usmanov, A.A.Okhunov, U.S.Salikhbaev and A.I.Vdovin // Phys. Part. Nucl. Lett. **7**(3), 185 (2010) [Pisma Fiz.Elem.Chastits At. Yadra **3**(159), 306 (2010)].
 - [8] M.Baglin // Nucl.Data Sheets, **96**, 611 (2002). // Nucl.Data Sheets, **77**(2), 125 (1996).
 - [9] B.Singh // Nucl.Data Sheets, **75**, 199 (1995).
 - [10] E.Browne, H.Junde // Nucl.Data Sheets, **87**, 15 (1999).
 - [11] S.M.Harris // Phys. Rev. **B138**, 509, (1965).
 - [12] A.A.Okhunov, Hasan Abu Kassim and Ph.N.Usmanov // Sains Malaysiana **40**(1), 1 (2011).
 - [13] R.B.Begzhanov, V.M.Belinkiy, I.I.Zalyubovskiy and A.B.Kuznichenko // "Handbook on Nuclear Physics", Tashkent, FAN, Vol.**1,2**, (1989).
 - [14] C.W.Reich, R.C.Greenwood, R.A.Lokken // Nucle. Phys. **A228**, 365 (1974).
 - [15] C.Fahlander, B.Varneestig, F.Backlin et al. // Nucl. Phys. **A147**, 157 (1992).
 - [16] A.Faessler, W.Greiner and R.K.Sheline // Nucl. Phys. **70**, 33 (1965).
 - [17] A.Arima and F.Iachello // Adv. Nucl. Phys. **13**, 139 (1984).
 - [18] R.F.Gasten, P.Von Brentano, W.R.Kane. // Phys. Rew. **C8**, 1035 (1973).




FIG. 2: (Color online) Comparison of the calculated and experimental energy spectra of positive-parity states for ^{170}Yb .




FIG. 3: (Color online) Comparison of the calculated and experimental energy spectra of positive-parity states for ^{172}Yb .




FIG. 4: (Color online) Comparison of the calculated and experimental energy spectra of positive-parity states for ^{174}Yb .




FIG. 5: (Color online) Structure of the wave-functions of 2_1^+ -band states for ^{170}Yb .




FIG. 6: (Color online) Structure of the wave-functions of 0_2^+ -band states for ^{170}Yb .




FIG. 7: (Color online) Structure of the wave-functions of 2_1^+ -band states for ^{172}Yb .




FIG. 8: (Color online) Structure of the wave-functions of 0_2^+ -band states for ^{172}Yb .




FIG. 9: (Color online) Structure of the wave-functions of 2_1^+ band states for ^{174}Yb .




FIG. 10: (Color online) Structure of the wave-functions of 0_2^+ band states for ^{174}Yb .